

# The redox couple of the cytochrome *c* cyanide complex: The contribution of heme iron ligation to the structural stability, chemical reactivity, and physiological behavior of horse cytochrome *c*

ABEL SCHEJTER,<sup>1</sup> MICHAEL D. RYAN,<sup>2</sup> ERICA R. BLIZZARD,<sup>3</sup> CHONGYAO ZHANG,<sup>3</sup> EMANUEL MARGOLIASH,<sup>4</sup> AND BENJAMIN A. FEINBERG<sup>3</sup>

<sup>1</sup>Sackler Institute of Molecular Medicine, Sackler Medical School, Tel-Aviv University, Tel-Aviv 69978, Israel

<sup>2</sup>Department of Chemistry, Marquette University, Milwaukee, Wisconsin 53233, USA

<sup>3</sup>Department of Chemistry, University of Wisconsin–Milwaukee, Milwaukee, Wisconsin 53211-3029, USA

<sup>4</sup>Department of Biochemistry, Molecular Biology, and Cell Biology, Northwestern University, Evanston, Illinois 60208, USA

(RECEIVED September 9, 2005; FINAL REVISION November 17, 2005; ACCEPTED November 17, 2005)


## Abstract

Contrary to most heme proteins, ferrous cytochrome *c* does not bind ligands such as cyanide and CO. In order to quantify this observation, the redox potential of the ferric/ferrous cytochrome *c*–cyanide redox couple was determined for the first time by cyclic voltammetry. Its  $E^{0'}$  was  $-240$  mV versus SHE, equivalent to  $-23.2$  kJ/mol. The entropy of reaction for the reduction of the cyanide complex was also determined. From a thermodynamic cycle that included this new value for the cyt *c* cyanide complex  $E^{0'}$ , the binding constant of cyanide to the reduced protein was estimated to be  $4.7 \times 10^{-3}$  LM<sup>-1</sup> or 13.4 kJ/mol (3.2 kcal/mol), which is 48.1 kJ/mol (11.5 kcal/mol) less favorable than the binding of cyanide to ferricytochrome *c*. For coordination of cyanide to ferrocyanochrome *c*, the entropy change was earlier experimentally evaluated as 92.4 Jmol<sup>-1</sup>K<sup>-1</sup> (22.1 e.u.) at 25 K, and the enthalpy change for the same net reaction was calculated to be 41.0 kJ/mol (9.8 kcal/mol). By taking these results into account, it was discovered that the major obstacle to cyanide coordination to ferrocyanochrome *c* is enthalpic, due to the greater compactness of the reduced molecule or, alternatively, to a lower rate of conformational fluctuation caused by solvation, electrostatic, and structural factors. The biophysical consequences of the large difference in the stabilities of the closed crevice structures are discussed.

**Keywords:** cytochrome *c*; cyanide coordination; cyt *c*–cyanide complex; thermodynamics; protein stability

In both the ferric and ferrous states, the heme iron of cytochrome *c* (cyt *c*) is hexacoordinated and lies at the center of a “closed heme coordination” structure

(George and Lyster 1958; Harbury et al. 1965) represented as



where the vertical lines signify the equatorial coordination by the heme pyrrole nitrogens. The iron has two axial ligands, Met<sub>80</sub> and His<sub>18</sub>. The His<sub>18</sub> imidazole is on what is commonly called the “right side” of the protein, and the


Reprint requests to: Abel Schejter, Sackler Institute of Molecular Medicine, Sackler Medical School, Tel-Aviv University, Tel-Aviv 69978, Israel; e-mail: molec03@post.tau.ac.il; fax: +972-3-547-2818.

Article and publication are at <http://www.proteinscience.org/cgi/doi/10.1110/ps.051825906>.

thioether sulfur of Met<sub>80</sub> is on the “left side” (Takano and Dickerson 1981).

Understanding the structural basis of the stability and the physiological behavior of this protein requires a thorough knowledge of the thermodynamic and kinetic consequences of this peculiar ligation of the metal. The Met<sub>80</sub>S–Fe<sup>3+</sup> bond is weaker than the Fe<sup>3+</sup>–ImidHis<sub>18</sub> bond, since the former can be broken by exogenous ligands, such as cyanide, which allows the measurement of its stability (George and Tsou 1952; George et al. 1967). However, an analogous evaluation of the same parameter in the ferrous state cannot be made, because typical ligands such as CO, O<sub>2</sub>, and CN<sup>−</sup> cannot break the Met<sub>80</sub>S–Fe<sup>2+</sup> bond. Indeed, since the earliest days of cyt *c* research it has been known that when the iron is in the reduced state, the protein is remarkably stable (Keilin 1930; Keilin and Hartree 1937), and, contrary to most heme proteins, unable to bind CO below pH 13 (Butt and Keilin 1962). This unusual characteristic of cyt *c* has, until now, remained only as a qualitative observation, lacking a definite quantitative measure and a consequent evaluation of its importance in the chemical and physiological behavior of the protein.


The redox-dependent ligand exchange of cyanide ion with Met<sub>80</sub>, proposed here to follow the 2×2 square mechanism (Evans 1990; Balducci and Costa 1993; Feinberg et al. 1998), is shown in Scheme 1. Since the redox potential of cytochrome *c* (see reaction C → D) (Scheme 1) and the affinity of its ferric state for cyanide



**Scheme 1.** The 2 × 2 square mechanism for the redox driven ligand exchange of cytochrome *c* in the presence of cyanide ion. Pr represents the native protein, and Pr\* symbolizes the protein after undergoing the structural changes concomitant with the displacement of the native ligand Met-80 away from the iron by CN<sup>−</sup>.

(reaction C → A) (Scheme 1) are known, it is possible, in principle, to provide a detailed thermodynamic characterization of the complete cycle by determining the redox potential of the cyanide complex (A → B) and deducing the stability of the Met<sub>80</sub>S–Fe<sup>2+</sup> bond (D → B) using the thermodynamic cycle shown in Scheme 1.

In Scheme 1, Pr represents the native protein, and Pr\* symbolizes the protein after undergoing the structural changes concomitant with the displacement of the native ligand (Met<sub>80</sub>) away from the iron by CN<sup>−</sup>. Once the redox potential of the cytochrome *c* cyanide complex is known (reaction A → B) (Scheme 1), the only unknown remaining in the thermodynamic cycle is the right-hand, bottom-to-top reaction (D → B, *k*<sub>f,BD</sub>), namely, the binding of cyanide to the ferrocycytochrome *c* iron:



It is known, however, that when the ferrocycytochrome *c* cyanide complex is formed at pH 13 by reducing the ferric cyanide complex, and then taken to neutrality, it dissociates readily with a half-life of 170 sec (George and Schejter 1964) back to Met<sub>80</sub>–Fe<sup>2+</sup>–His<sub>18</sub>. This has precluded an accurate measurement of the equilibrium reduction potential of this complex by conventional equilibrium titration techniques, such as potentiometry or spectroelectrochemistry.

Cyclic voltammetry (CV) (Bard and Faulkner 2001), in combination with modern computer simulation methods, permits the determination of the redox potential of species with short half-lives, such as the CN–Fe<sup>3+/2+</sup>–His<sub>18</sub> redox couple. We report in this paper the indirect measurement of the stability of the Met<sub>80</sub>S–Fe<sup>2+</sup> bond of cytochrome *c* using CV. Our results show that the Met<sub>80</sub>S–Fe<sup>2+</sup> bond is 48.1 kJ/mol (11.5 kcal/mol) more stable (ΔΔ*G*) in the reduced state than in the oxidized state, thus, providing the first quantitative answer to a fundamental and long-standing problem in the chemistry of this paradigmatic heme protein. On this basis, we are able to explain the large difference in the stabilities of the two oxidation states of the protein against denaturation. We also note how this difference plays a major role in the regulation of the redox potential of cytochrome *c* and in protecting the reduced iron from being oxidized by oxygen.

## Materials and methods

### Preparative procedures

Horse heart cytochrome *c* was prepared as described by Margoliash and Walassek (1967) and Brautigan et al. (1978).

*Electrochemical methods**Cyclic voltammetry*

While CV is not as sensitive as square-wave voltammetry (SWV), it possesses some important advantages, including (1) the direct observation of the redox couples and the relative contributions of reducing/oxidizing current to each couple and (2) a more direct interpretation of the voltammogram, especially as a function of the scan rate. Cyclic voltammetric data at two or more scan rates are typically simulated within the framework of a specific mechanism (e.g., Scheme 1), and can provide the  $E^{0'}$  of the participating redox couples and/or the associated equilibrium and rate constants. In CV, a triangular waveform is applied to the electrode (Bard and Faulkner 2001). The potential is first scanned in one direction and then, at some point, switched and scanned in the reverse direction. For example, for a ferricyt *c* solution when the potential is scanned in the negative direction, the ferricyt *c* is reduced to the ferrous form as the potential approaches and goes past its  $E^{0'}$ ; a cathodic peak current is observed; and then, upon scanning the potential in the reverse direction, the reduced form is reoxidized to give the anodic peak current. The complete voltammogram gives a pair of peaks of the same current amplitude but in the opposite direction with the  $E^{0'}$  being between the two peak potentials with a peak potential separation of  $60 \text{ mV}/n$  ( $n$  = number of electrons per equivalent used in the redox reaction) when the redox reaction is fast and reversible (Nernstian).


*Simulation of cyclic voltammograms*

A relatively new and fast method for simulating cyclic voltammograms has been developed (Rudolph 1990, 1995; Rudolph et al. 1994), which, in principle, permits the determination of the  $E^{0'}$ , equilibrium constants and rates of production of new products and intermediates in the context of a specific mechanism and as a function of scan rate. The simulation program has now been formalized as the computer software, Digisim (Bioanalytical Systems, Inc.), which is used in this work.

*Square-wave voltammetry*

SWV is a voltammetric method that provides a high signal to noise response (Ramely and Kranse 1969). In SWV a potential is applied over time in the form of a square wave superimposed on a staircase. The applied potential is progressively stepped in fixed increments and pulsed positively and negatively at each step with a specific step potential, the square-wave amplitude. Experimentally, the positive (forward) and negative (reverse) currents are collected separately and then the differences are taken for forward and reverse currents to provide the observed "net" SW voltammogram. In the

absence of kinetic complications, the potential at which the voltammetric peak is observed defines the equilibrium reduction potential,  $E^{0'}$ . In this paper, the variation of  $E^{0'}$  with temperature was obtained with SWV and used to determine the entropy of reaction,  $\Delta S^{0'}_{\text{Rxn}}$ , for reduction reaction:



$\Delta S^{0'}_{\text{Rxn}}$  was determined via nonisothermal electrochemical measurements of the equilibrium reduction potential,  $E^{0'}$ , as a function of temperature. Error analysis for the experiments in Figure 2 and shown in Table 1 were obtained with weighted least-squares method to calculate  $\Delta S$  and  $\Delta H$ . It was assumed that the potentials observed were  $\pm 1.5 \text{ mV}$ . The experimental details for using SWV to determine the  $\Delta S^{0'}_{\text{Rxn}}$  have been reported (Bertrand et al. 1995; Feinberg et al. 1998). In earlier work, SWV and CV were used to understand the redox-driven axial ligand exchange of the Phe82His variant of iso-1-cytochrome *c* (Feinberg et al. 1998), which followed the  $2 \times 2$  square mechanism similar to that shown in Scheme 1.

*Instrumentation*

All electrochemical experiments were done with a BAS 100 B/W Electrochemical Analyzer (Bioanalytical Systems). Potentials were reported against either the Standard Hydrogen Electrode (SHE) or the Ag/AgCl/3 M KCl reference electrode. To convert Ag/AgCl potentials to the SHE, 210 mV was added to that potential. A 0.1 M solution of KCN buffered at pH 7.0 with Tris was prepared. Samples (100  $\mu\text{L}$ ) of horse cytochrome *c* were then microdialyzed into this cyanide solution overnight at  $4^\circ\text{C}$  and used immediately thereafter. Triple-distilled water (Purification Systems) with a conductivity of 18 Mohms/cm was used throughout. Final protein concentrations were 0.5–1 mM. All experiments were done at a bis(4-pyridyl) disulfide modified gold electrode. Experimental details, including the modification of the gold electrode with 5 mM bis(4-pyridyl) disulfide (Aldrich

**Table 1.** Thermodynamic parameters for the reduction of horse cytochrome *c* and its cyanide complex pH 7.0 and  $25^\circ\text{C}$

	$E^{0'}$ mV	$\Delta G$ kJ/mol kcal/mol	$\Delta H$ kJ/mol kcal/mol	$\Delta S$ J/mol-K e.u.
Cytochrome <i>c</i> <sup>a</sup>	+260	-25.1 -6.0	-69.9 -16.7	-150.6 -36.0
Cytochrome <i>c</i> cyanide	-240	+23.2 $\pm$ 0.2 +5.5 $\pm$ 0.03	-3.8 $\pm$ 1 -0.9 $\pm$ 0.3	-90.0 $\pm$ 4 -21.5 $\pm$ 1

<sup>a</sup>George and Lyster 1958; Margalit and Schejter 1973.


Chemical Co.) have been described (Feinberg et al. 1998).

A specially designed anaerobic semimicrocell that permitted samples of  $\sim 30\text{-}\mu\text{L}$  volume to be examined was used for the electrochemical experiments (Smith and Feinberg 1990; Smith et al. 1991). Oxygen was removed by passing wet 99.99% pure argon over the sample and through the cell for 20 min prior to the experiment. The reference electrode, which was separated from the sample with a double frit, was the Ag/AgCl/3 M KCl electrode (Bioanalytical Systems). The counter electrode was 18-gauge platinum wire. For the entropy of reaction measurements,  $E^0$ 's were measured at 1.6°, 9.4°, 18.0°, 28.1°, and 37.1°C.

## Results

### Cyclic voltammetry

Figure 1 shows the cyclic voltammogram of cytochrome *c* in 0.1 M KCN (pH 7.0), 25°C, at a 50 mV/sec scan rate, along with the computer simulation of that voltammogram with Digisim using the  $2 \times 2$  square mechanism shown in Scheme 1. In following the course of the voltammogram (see Fig. 1; Scheme 1), when the voltammetric experiment was started at 0.3 V, all the cytochrome near the electrode surface was oxidized. As the applied potential was moved in the negative (reducing) direction, a small cathodic peak (C  $\rightarrow$  D) close to the  $E^0$  of native horse cytochrome *c* was observed. This peak, assigned to the reduction of the ferric cyt *c*, is quite small and shows that nearly all the cyt *c* has the Met<sub>80</sub> ligand displaced by CN<sup>-</sup>. That is, the cytochrome is mostly in the form CN—Fe<sup>3+</sup>—His<sub>18</sub>, leaving very little in the Met<sub>80</sub>—Fe<sup>3+</sup>—His<sub>18</sub> form. As the scan was




**Figure 1.** Cyclic voltammogram of 1 mM horse cytochrome *c* in the presence of 0.1 M KCN at pH 7.0, Tris buffer, 25°C at a scan rate of 50 mV/sec. The circles (O) are the experimental data; the solid line, the computer simulation of the CV with Digisim using the  $2 \times 2$  mechanism shown in Scheme 1.

continued in the negative direction, a much larger cathodic (reduction) peak (A  $\rightarrow$  B) was observed at  $\sim -500$  mV (vs. Ag/AgCl), which is assigned to the reduction of the ferric cyt *c* cyanide complex. At  $-600$  mV, the direction of the potential scan was switched and moved in the positive (oxidizing) direction. Soon after the return scan was initiated, a nearly full “reverse current” peak (B  $\rightarrow$  A) was observed for the reoxidation of the ferrous cyt *c* cyanide complex. This indicated that the cyanide was still the primary sixth axial ligand, even with the iron in the ferrous state. The reason that this large “reverse current” was observed is that on the timescale of the voltammetric experiments, the rate of dissociation of the ferrocyanide is slow, so that most of the cyt *c* was still in the CN<sup>-</sup> coordinated form (B) at the time it was reoxidized (A). As the scan was continued in the oxidizing direction, another small peak was observed near the redox potential of the native cyt *c* (D  $\rightarrow$  C). Summing up, in the complete voltammogram two relatively undistorted yet interconnected (see Scheme 1) redox couples were observed: one for the native cyt *c* and one for the cyt *c*–cyanide complex. Fortunately, two factors combine to allow us to observe the  $E^0$  of cyt *c*–CN complex: one is the large difference between the  $E^0$  of the cyt *c* and cyt *c*–CN complex redox couples, i.e., 260 mV versus  $\sim -220$  mV, respectively, and the other is the slow rate of the ferrous cyt *c*–cyanide complex dissociation relative to the potential scan rate. Thus, even prior to simulation, which takes into account the kinetic effects, the voltammogram in Figure 1 provides an approximate value for the  $E^0 \approx -220$  mV versus SHE for the CN—Fe<sup>3+/2+</sup>—His<sub>18</sub> couple and  $E^0 = +260$  mV for the native cyt *c*.


### Simulation of the cyclic voltammograms: Determining the $E^0$ of the cytochrome *c* cyanide complex via the $2 \times 2$ square mechanism

Shown in Figure 1 is the voltammogram and its simulation of cyt *c* in 0.1 M NaCN at a scan rate of 50 mV/sec. The other scan rate used was 100 mV/sec and was also used in the simulation. Simulations made with Digisim using the  $2 \times 2$  square mechanism (Scheme 1) took into account the extent to which kinetics shifted the observed reduction potential. In fact, kinetic effects are real but not extensive at the scan rates used here, and the cyclic voltammogram is dominated by equilibrium processes. In starting the simulation, the approximate  $K_{\text{eq}} = k_{f,AC}/k_{b,AC} \approx 0.01$  used. This approximate  $K_{\text{eq}}$  was obtained from the ratio of the cathodic voltammetric currents of peaks C  $\rightarrow$  D and A  $\rightarrow$  B of the voltammogram. The relatively fast heterogenous electron

transfer rate constant of  $k_s = 3 \times 10^{-3}$  cm/sec was employed since it gave the best fit for both scan rates and was used in conjunction with the cross-reaction:



where  $K_{\text{cross-rxn}}$  was quite large by virtue of the relative  $E^0$ 's of the wild-type cyt *c* (+260 mV) and its cyanide complex (-240 mV). For the equilibrium,



D

B

the simulated (calculated) value of  $K_{\text{eq}} = K_{\text{BD}} = k_{f,\text{BD}}/k_{b,\text{BD}}$  was  $5.2 \times 10^{-7}$ .

The scan rates employed to obtain an accurate  $E^0$  (50 and 100 mV/sec) were relatively fast on the timescale of the dissociation of  $\text{CN-Fe}^{2+}\text{-His}_{18}$ , and thus did not place these voltammograms well into the kinetic region, i.e., where significant potential shifts and changes in cathodic and anodic currents would have been observed for both the  $\text{Met}_{80}\text{-Fe}^{3+/2+}\text{-His}_{18}$  and  $\text{CN-Fe}^{3+/2+}\text{-His}_{18}$  redox couples. Moreover, by choosing the somewhat faster scan rates (where the voltammograms were in the "equilibrium region"), we have obtained an accurate value for the  $E^0$  of the  $\text{CN-Fe}^{3+/2+}\text{-His}_{18}$  couple, but obtained less precise information on the kinetic parameters shown in Scheme 1. To explore the kinetics of the redox driven cyanide ligand exchange under these conditions, voltammetric experiments would need to be done at significantly slower scan rates.


#### Square-wave voltammetry as a function of temperature

Figure 2 shows the variation of the reduction potential of the  $\text{CN-Fe}^{3+/2+}\text{-His}_{18}$  couple as a function of temperature, which in turn, provided the entropy of reaction (Feinberg et al. 1998) used to calculate the overall entropy change (Taniguchi et al. 1980) via the expression:  $\Delta S^0 = \Delta S^0_{\text{Rxn}} - 66.5 \text{ J/mol} \cdot \text{K}$ .

#### Calculation of the thermodynamics of cyanide binding

From the cycle of Scheme 1, it follows, starting with native cyt *c*, that

$$\begin{aligned} \Delta G(c^{3+} \rightarrow c^{2+}) + \Delta G(c^{2+} \rightarrow \text{CN-}c^{2+}) + \\ \Delta G(\text{CN-}c^{2+} \rightarrow \text{CN-}c^{3+}) + \\ \Delta G(\text{CN-}c^{3+} \rightarrow c^{3+}) = 0 \end{aligned} \quad (1)$$



**Figure 2.** Temperature dependence of the equilibrium reduction potential ( $E^0$ ) of the cyanide complex of horse cytochrome *c*.

where all the  $\Delta G$  terms are written in a counterclockwise direction as one moves around Scheme 1. The bottom reaction (C  $\rightarrow$  D) is the reduction of ferricyt *c* ( $E^0 = +260$  mV) to ferrocyt *c* for which  $\Delta G(c^{3+} \rightarrow c^{2+}) = -25.1$  kJ/mol (6.0 kcal/mol) (Margalit and Schejter 1973). The top reaction (B  $\rightarrow$  A), from right to left, is the oxidation of ferrous cyt *c* cyanide to ferric cyt *c* cyanide,  $\Delta G(\text{CN-}c^{2+} \rightarrow \text{CN-}c^{3+})$ ,  $E^0 = -0.240$  mV) which corresponds to  $-23.0$  kJ/mol (5.5 kcal/mol). The left top-to-bottom reaction, A  $\rightarrow$  C,  $\Delta G(\text{CN-}c^{3+} \rightarrow c^{3+})$ , is the dissociation of the cyanide from the ferric protein, for which the free energy change is  $+34.7$  kJ/mol (8.3 kcal/mol) (George and Tsou 1952; George et al. 1967). When these values are introduced into Equation 1, the free energy of binding of cyanide to ferrocyt *c*, (D  $\rightarrow$  B),  $\Delta G(\text{CN}^- + c^{2+} \rightarrow \text{CN-}c^{2+})$  at 25°C was calculated to be 13.4 kJ/mol (3.2 kcal/mol). This is equivalent to a binding constant of  $4.7 \times 10^{-3} \text{ LM}^{-1}$ , a distinctly low affinity that explains why the cyanide complex of reduced cyt *c* cannot be formed directly at neutral pH.

Applying our electrochemical data (see Fig. 2) to the equation  $d(\Delta G)/dT = \Delta S$ , the entropy change attending the reduction of the ferricyt *c*-cyanide complex was evaluated as  $-90 \text{ J/mol} \cdot \text{K}$  (21.5 e.u.) at 25°C. Expanding the free energy terms into their enthalpic and entropic components, an equation similar to Equation 1 can be written for the entropies of the reactions described in Scheme 1. Since the entropies of reduction and cyanide binding of ferricyt *c* are known to be  $-150.6 \text{ J/mol} \cdot \text{K}$  (36 e.u.) (Margalit and Schejter 1973) and  $132 \text{ J/mol} \cdot \text{K}$  (31.5 e.u.) (George and Tsou 1952; George et al. 1967), respectively, the entropy of cyanide binding to ferrocyt *c* was evaluated as  $92.4 \text{ J/mol} \cdot \text{K}$  (22.1 e.u.).

These results are listed in Tables 1 and 2, together with the corresponding heats of binding of cyanide to ferric and ferrous cyt *c*. The latter was evaluated, using the Gibbs equation  $\Delta G = \Delta H - T \cdot \Delta S$ , as  $41.0$  kJ/mol (9.8 kcal/mol).

## Discussion

The preceding results show that the free energy change (difference) concomitant with the binding of cyanide to the ferrous heme iron of cyt *c* is 48.1 kJ/mol (11.5 kcal/mol) less favorable than when the iron is in the ferric state (see Table 2). The data in Table 2 show that ~75% of the difference between the stabilities ( $\Delta\Delta G$ ) of the closed heme coordination structures (George and Lyster 1958; Takano and Dickerson 1981) of the ferric and ferrous states arises from the difference between the enthalpies ( $\Delta\Delta H$ ) of the reactions of the latter, which amounts to 36.4 kJ/mol (8.7 kcal/mol), i.e., 36.4/48.1 kJ/mol  $\approx$  0.75. The term “heme coordination stability” is synonymous with the earlier term “heme crevice stability,” i.e., the amount of free energy required to break the methionyl S–Fe bond. The fact that the driving force of cyanide binding to ferric cyt *c* has a large favorable entropy change, led to earlier speculations that attributed the lack of reactivity of the ferrous form to the absence of an entropic contribution (George and Schejter 1964; Margoliash and Schejter 1966). Indeed, cyanide coordination to ferrous cyt *c* does make an entropy contribution ( $T \cdot \Delta\Delta S$ ) to the free energy of the binding reaction, but of only ~40 J/mol·K (9.4 e.u.), which amounts to 11.8 kJ/mol (2.8 kcal/mol) at 25°C (see Table 2), or ~25% of the total free energy.

The large difference in enthalpies can be accounted for by two major factors: electrostatic and structural. First, an important contribution must stem from the fact that cyanide binding to the ferric state implies a cancellation of a charge that is absent in the ferrous state. The second factor should involve the structural differences between the ferric and ferrous proteins (Takano and Dickerson 1981) in the region of the molecule to which the Met<sub>80</sub> side chain is necessarily displaced upon breaking its coordination to the iron. Although structural studies

indicate that packing of the reduced and the oxidized proteins is essentially the same in both oxidation states (Brayer and Murphy 1996), there is significant evidence that the oxidized and the reduced molecules differ in their hydrodynamic properties. Thus, the intrinsic viscosity of the reduced state is lower than that of the oxidized state (Fisher et al. 1973), and the radius of gyration (Trehwella et al. 1988) and adiabatic compressibility (Eden et al. 1982) are correspondingly smaller. All these differences indicate a dynamically more “open” structure in the oxidized state and a more compact structure in the reduced state. In the now more compact ferrous cyt *c*, it is more difficult to find space for the displaced Met<sub>80</sub> side chain, further impeding CN<sup>−</sup> binding. This difference must also play a significant role in the regulation of the redox potential of cyt *c* by virtue of stabilizing the reduced form. Further work will be performed to parse the diverse contributions to this parameter arising from intrinsic ligand-metal affinities, protein conformation differences, and overall electrostatic influences.

Moreover, the preceding results also explain, quantitatively, the extreme resistance of the ferrous cyt *c* coordination structure being broken (i.e., breaking the axial methionyl S–Fe<sup>2+</sup> heme coordination bond) by exogenous ligands including oxygen, and thus its resistance to oxidation by oxygen. Another demonstration that reduction increases the affinity of the heme iron for a methionyl sulfur is the fact that substitution of histidine for the Phe<sub>82</sub> in iso-1 cyt *c* results in the displacement of Met<sub>80</sub> from the ferric iron by the new residue (Hawkins et al. 1994). However, upon reduction of the metal, the normal Met<sub>80</sub>–Fe<sup>2+</sup> bond is restored (Schejter et al. 1996; Feinberg et al. 1998). For comparison, in Met-65,80-dicarboxymethyl cyt *c*, a chemically modified cyt *c* in which the alkylated Met<sub>80</sub> sulfur is displaced from the iron, the reduced metal binds oxygen rapidly and is oxidized to the ferric state within a few minutes (Schejter and Aviram 1970). This ability of native cyt *c* to remain reduced in the presence of oxygen, which is essential for its proper function in the respiratory chain, depends directly on the high stability of the native heme crevice.

It is interesting that the high stability of the reduced cyt *c* closed-crevice structure is reflected almost quantitatively in the increased overall stability of the protein in the reduced state. Specifically, the free energy of chemical denaturation of horse ferrous cytochrome *c* is 77.0 kJ/mol (18.4 kcal/mol) (Jones et al. 1993), while the corresponding value for ferric cyt *c* is 33.9 kJ/mol (8.1 kcal/mol) (Knapp and Pace 1974; Schejter et al. 1998). Strikingly, this higher stability of the reduced protein structure of 10.3

**Table 2.** Thermodynamic parameters for the reactions of ferric and ferrous cytochrome *c* with cyanide, at pH 7.0 and 25°C<sup>a</sup>

	$\Delta G$ kJ/mol kcal/mol	$\Delta H$ kJ/mol kcal/mol	$\Delta S$ J/mol·K e.u.	$T\Delta S$ kJ/mol
Ferric cytochrome <i>c</i>	−34.7	4.6	132.0	39.3
	−8.3	1.1	31.5	
Ferrous cytochrome <i>c</i>	13.4	41.0	92.4	27.5
	3.2	9.8	22.1	

Note:  $\Delta\Delta G = 48.1$  kJ/mol (11.5 kcal/mol);  $\Delta\Delta H = 36.4$  kJ/mol;  $T(\Delta\Delta S) = 11.8$  kJ/mol

<sup>a</sup> George and Tsou 1952; George 1967; Margalit and Schejter 1973.

kcal/mol (43.1 kJ/mol) is very close to the increase in stability of the native heme crevice upon reduction, i.e., 48.1 kJ/mol (11.5 kcal/mol;  $\Delta\Delta G$  from Table 2). This correspondence of free energies may very well be coincidental, since it has been shown with *Saccharomyces cerevisiae* iso-1-cyt *c* that the source of the increase in thermal stability of the reduced the protein is nearly entirely entropic (Cohen and Pielak 1995), albeit at pH 4.6. In rationalizing the increased enthalpic stabilization of heme coordination (a local effect) versus the entropically driven increase in overall stability of the reduced protein, it is important to point out that the energetics presented in this paper refer specifically to the enthalpic barrier to breaking the Met<sub>80</sub>-Fe<sup>2+</sup> bond by both endogenous ligands and exogenous ligands such as cyanide, imidazole, or oxygen. Since there is evidence that reduction and increased overall stability are concomitant with the protein becoming more compact (see the above discussion), the more compact structure may make ligand exchange energetically more difficult just from simple molecular mechanical considerations. Invoking increased Met<sub>80</sub>-Fe<sup>2+</sup> bond strength and thus an increased enthalpic contribution is not a likely option, since, again, in the thermal denaturation study of the overall stability of reduced iso-1 cyt *c*, Cohen and Pielak (1995) found that the enthalpic change upon reduction was destabilizing.

In summary, the preceding results provide a quantitative measure of the difference in the stabilities of the closed structures of reduced and oxidized cyt *c*. They also demonstrate that this feature plays an important role in the determination of the redox potential of this heme protein, in its resistance to oxidation by molecular oxygen, and on the stabilization of the native protein structure.

### Acknowledgments

We acknowledge support of this research by a National Institutes of Health Grant GM 55892 to B.A.F. and a grant from the Edward Mallinckrodt Jr. Foundation, St. Louis, MO, to E.M.

### References

- Balducci, G. and Costa, G. 1993. The four-member square scheme in cyclic voltammetry: General solution for Nerstian electron transfers. *J. Electroanal. Chem.* **348**: 355–365.
- Bard, A.J. and Faulkner, L.R. 2001. *Electrochemical methods: Fundamentals and applications*. Wiley & Sons, Inc., New York.
- Bertrand, P., Mbarki, O., Asso, M., Blanchard, L., Guerlesquin, F., and Tegoni, M. 1995. Control of the redox potential in *c*-type cytochromes: Importance of the entropic contribution. *Biochemistry* **34**: 11071–11079.
- Brautigan, D.L., Ferguson-Miller, S., and Margoliash, E. 1978. Mitochondrial cytochrome *c*: Preparation and activity of native and chemically modified cytochromes *c*. *Methods Enzymol.* **53**: 128–164.
- Brayer, G.D. and Murphy, M.E.P. 1996. Structural studies of eukaryotic cytochromes *c*. In *Cytochrome c—A multidisciplinary approach* (eds. R.A. Scott and A. Grant Mauk), pp. 103–166. University Science Books, Sausalito, CA.
- Butt, J.N. and Keilin, D. 1962. Absorption spectra and some other properties of cytochrome *c* and of its compounds with ligands. *Proc. R. Soc. Lond. Ser. B* **156**: 429–458.
- Cohen, D.S. and Pielak, J.G. 1995. Entropic stabilization of cytochrome *c* upon reduction. *J. Am. Chem. Soc.* **117**: 1675–1677.
- Eden, D., Matthew, J.B., Rosa, J.J., and Richards, F.M. 1982. Increase in apparent compressibility of cytochrome *c* upon oxidation. *Proc. Natl. Acad. Sci.* **79**: 815–819.
- Evans, D.H. 1990. Solution electron-transfer reactions in organic and organometallic electrochemistry. *Chem. Rev.* **90**: 739–751.
- Feinberg, B.A., Liu, X., Ryan, M.D., Schejter, A., Zhang, C., and Margoliash, E. 1998. Direct voltammetric observation of redox driven changes in axial coordination and intramolecular rearrangement of the phenylalanine-82-histidine variant of yeast iso-1-cytochrome *c*. *Biochemistry* **37**: 13091–13101.
- Fisher, W.R., Taniuchi, H., and Anfinsen, C.B. 1973. On the role of heme in the formation of the structure of cytochrome *c*. *J. Biol. Chem.* **248**: 3188–3195.
- George, P. and Lyster, R.L.J. 1958. Crevice structures in hemoprotein reactions. *Proc. Natl. Acad. Sci.* **44**: 1013–1029.
- George, P. and Schejter, A. 1964. The reactivity of ferrocycytochrome *c* with iron-binding ligands. *J. Biol. Chem.* **239**: 1504–1508.
- George, P. and Tsou, C.L. 1952. The reaction between hydrocyanic acid, cyanide ion and ferricytochrome *c*. *Biochemistry* **50**: 440–452.
- George, P., Glauser, S.C., and Schejter, A. 1967. The reactivity of ferricytochrome *c* with ionic ligands. *J. Biol. Chem.* **242**: 1690–1695.
- Harbury, H.A., Cronin, Y.R., Fanger, M.W., Hettlinger, T.P., Murphy, A.J., Myer, Y.P., and Vinogradov, S.N. 1965. Complex formation between methionine and a heme peptide from cytochrome *c*. *Proc. Natl. Acad. Sci.* **54**: 1658–1664.
- Hawkins, B.K., Hilgenwillis, S., Pielak, G.J., and Dawson, J.H. 1994. Novel axial ligand interchange in Cytochrome *c*: Incorporation of a histidine at position 82 leads to displacement of the wild-type methionine-80 ligand. *J. Am. Chem. Soc.* **116**: 3111–3112.
- Jones, C.M., Henry, E.R., Hu, Y., Chan, C.-K., Luck, S.D., Bhuyan, A., Roder, H., Hofrichter, J., and Eaton, W.A. 1993. Fast events in protein folding initiated by nanosecond laser photolysis. *Proc. Natl. Acad. Sci.* **90**: 11860–11864.
- Keilin, D. 1930. Cytochrome and intracellular oxidase. *Proc. R. Soc. Lond. B* **106**: 418–444.
- Keilin, D. and Hartree, E.F. 1937. Preparation of pure cytochrome *c* from heart muscle and some of its properties. *Proc. R. Soc. Lond. B* **122**: 298–313.
- Knapp, J.A. and Pace, C.N. 1974. Guanidine hydrochloride and acid denaturation of horse, cow and *Candida krusei* cytochromes *c*. *Biochemistry* **13**: 1289–1294.
- Margalit, R. and Schejter, A. 1973. Cytochrome *c*: Thermodynamic study of the relations between oxidation state, ion binding, and structural parameters. I. Effects of temperature, pH, and electrostatic media on the standard redox potential. *Eur. J. Biochem.* **32**: 492–499.
- Margoliash, E. and Schejter, A. 1966. Cytochrome *c*. *Adv. Protein Chem.* **21**: 126–213.
- Margoliash, E. and Walassek, O.F. 1967. Cytochrome *c* from vertebrate and invertebrate sources. *Methods Enzymol.* **10**: 339–348.
- Ramely, L. and Kranse Jr., M.S. 1969. Theory of square wave voltammetry. *Anal. Chem.* **41**: 1362–1365.
- Rudolph, M. 1990. An algorithm of general application for the digital simulation of electrochemical processes. *J. Electroanal. Chem.* **292**: 1–7.
- . 1995. Digital simulations with the fast implicit finite difference algorithm: The development of a general simulator for electrochemical processes. In *Physical electrochemistry: Principles, methods, and applications* (ed. I. Rubinstein), pp. 81–129. Marcel Dekker, Inc., New York.
- Rudolph, M., Reddy, D.P., and Feldberg, S.W. 1994. A simulator for cyclic voltammetric responses. *Anal. Chem.* **66**: A589–A600.
- Schejter, A. and Aviram, I. 1970. The effects of alkylation of methionyl residues on the properties of horse cytochrome *c*. *J. Biol. Chem.* **295**: 1552–1557.
- Schejter, A., Taler, G., Navon, G., Liu, X.-J., and Margoliash, E. 1996. Oxidation state-induced change of iron ligand in the phenylalanine-82

- to histidine mutant of yeast iso-1-cytochrome *c*. *J. Am. Chem. Soc.* **118**: 477–478.
- Schejter, A., Sanishvili, R., Qin, W., and Margoliash, E. 1998. A comparison of local and global effects of mutations of mammalian cytochrome *c*. *Chemtracts—Biochem. Mol. Biol.* **11**: 713–728.
- Smith, E.T. and Feinberg, B.A. 1990. Redox properties of several bacterial ferredoxins using square wave voltammetry. *J. Biol. Chem.* **265**: 14371–14376.
- Smith, E.T., Bennett, D.W., and Feinberg, B.A. 1991. Redox properties of 2[4Fe-4S] ferredoxins. *Anal. Chim. Acta* **251**: 27–33.
- Takano, T. and Dickerson, R.E. 1981. Conformation change of cytochrome *c*. I. Ferrocycytochrome structure refined at 1.5 Å resolution. *J. Mol. Biol.* **153**: 74–94.
- Taniguchi, V., Sailasuta-Scott, N., Anson, F.C., and Gray, H.B. 1980. Thermodynamics of metalloprotein electron transfer reactions. *Pure Appl. Chem.* **52**: 2275–2281.
- Trewhella, J., Carlson, V.A., Curtis, E.H., and Heidorn, D.B. 1988. Differences in the solution structure of oxidized and reduced cytochrome *c* measured by small-angle X-ray scattering. *Biochemistry* **27**: 1121–1125.